

Chapter 4 Anxiety

Adams TB, Moore MT, Dye J. The relationship between physical activity and mental health in a national sample of college females. *Women Health*. 2007;45(1):69-85.

Asikainen TM, Miilunpalo S, Oja P, Rinne M, Pasanen M, Uusi-Rasi K, Vuori I. Randomised, controlled walking trials in postmenopausal women: the minimum dose to improve aerobic fitness? *Br J Sports Med*. 2002 Jun;36(3):189-94

Audette JF, Jin YS, Newcomer R, Stein L, Duncan G, Frontera WR. Tai Chi versus brisk walking in elderly women. *Age Ageing*. 2006 Jul;35(4):388-93. Epub 2006 Apr 19.

Bartholomew JB, Linder DE. State anxiety following resistance exercise: the role of gender and exercise intensity. *J Behav Med*. 1998 Apr;21(2):205-19.

Bartholomew JB, Morrison D, Ciccolo JT. Effects of acute exercise on mood and well-being in patients with major depressive disorder. *Med Sci Sports Exerc*. 2005 Dec;37(12):2032-7

Baruch DE, Swain RA, Helmstetter FJ. Effects of exercise on Pavlovian fear conditioning. *Behav Neurosci*. 2004 Oct;118(5):1123-7

Bhui, K., & Fletcher, A. (2000). Common mood and anxiety states: Gender differences in the protective effects of physical activity. *Social Psychiatry and Psychiatric Epidemiology*, 35, 28-35.

Binder E, Droste SK, Ohl F, Reul JM. Regular voluntary exercise reduces anxiety-related behaviour and impulsiveness in mice. *Behav Brain Res*. 2004 Dec 6;155(2):197-206.

Broman-Fulks JJ, Berman ME, Rabian BA, Webster MJ. Effects of aerobic exercise on anxiety sensitivity. *Behav Res Ther*. 2004 Feb;42(2):125-36.

Bonhauer M, Fernandez G, Püschel K, Yañez F, Montero J, Thompson B, Coronado G. Improving physical fitness and emotional well-being in adolescents of low socioeconomic status in Chile: results of a school-based controlled trial. *Health Promot Int*. 2005 Jun;20(2):113-22.

Breus, M. J., & O'Connor, P. J. (1998). Exercise-induced anxiolysis: A test of the "time out" hypothesis in high anxious females. *Medicine and Science in Sports and Exercise*, 30, 1107-1112.

Broocks A, Bandelow B, Pekrun G, George A, Meyer T, Bartmann U, Hillmer-Vogel U, Rüther E. Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder. *Am J Psychiatry*. 1998 May;155(5):603-9

Broocks A, Meyer T, Opitz M, Bartmann U, Hillmer-Vogel U, George A, Pekrun G, Wedekind D, Rüther E, Bandelow B. 5-HT1A responsivity in patients with panic disorder before and after

treatment with aerobic exercise, clomipramine or placebo. *Eur Neuropsychopharmacol.* 2003 May;13(3):153-644

Chaouloff, F., Laude, D. & Elghozi, J. (1989). Physical exercise: evidence for differential consequences of tryptophan on 5-HT synthesis and metabolism in central serotonergic cell bodies and terminals. *Journal of Neural Transmission* (Gen Sect) 78, 121-130.

Chaouloff, F. (1993). Physiopharmacological interactions between stress hormones and central serotonergic systems. *Brain Res. Rev.*, 18, 1-32.

Chaouloff F (1997) Effects of acute physical exercise on central serotonergic systems. *Medicine and Science in Sports and Exercise* 29(1): 58-62.

Chow YW, Tsang HW. Biopsychosocial effects of qigong as a mindful exercise for people with anxiety disorders: a speculative review. *J Altern Complement Med.* 2007 Oct;13(8):831-9.

Cox, R.H., Thomas, T.R. and Davis, J.E. (2000) Delayed anxiolytic effect associated with an acute bout of aerobic exercise. *Journal of Exercise Physiology Online* 3, 59-66.

Cox RH, Thomas TR, Hinton PS, Donahue OM. Effects of acute 60 and 80% VO₂max bouts of aerobic exercise on state anxiety of women of different age groups across time. *Res Q Exerc Sport.* 2004 Jun;75(2):165-75

De Moor MH, Beem AL, Stubbe JH, Boomsma DI, De Geus EJ. Regular exercise, anxiety, depression and personality: a population-based study. *Prev Med.* 2006 Apr;42(4):273-9. Epub 2006 Jan 24.

DeVries, H. A. (1981). Tranquilizer effect of exercise: A critical review. *The Physician and Sports Medicine*, 9, 46–55.

DeVries, H. A., Wiswell, R. A., Bulbulian, R., & Moritani, T. (1981). Tranquilizer effect of exercise: Acute effects of moderate aerobic exercise on spinal reflex activation level. *American Journal of Physical Medicine*, 60, 57–

DiLorenzo TM, Bargman EP, Stucky-Ropp R, Brassington GS, Frensch PA, LaFontaine T. Long-term effects of aerobic exercise on psychological outcomes. *Prev Med* 1999;28:75– 85.

DiLorenzo TM, Bargman EP, Stucky-Ropp R, Brassington GS, Frensch PA, LaFontaine T. Long-term effects of aerobic exercise on psychological outcomes. *Prev Med.* 1999 Jan;28(1):75-85.

Dishman RK (1997) Brain monoamines, exercise and behavioral stress: animal models. *Medicine and Science in Sports and Exercise*, 29(1):63-74.

Droste SK, Gesing A, Ulbricht S, Müller MB, Linthorst AC, Reul JM. Effects of long-term voluntary exercise on the mouse hypothalamic-pituitary-adrenocortical axis. *Endocrinology*. 2003 Jul;144(7):3012-23

Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. The DOSE study: a clinical trial to examine efficacy and dose response of exercise as treatment for depression. *Control Clinical Trials*. 2002 Oct;23(5):584-603.

Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc*. 2001 Jun;33(6 Suppl):S587-97; discussion 609-10.

Esquivel G, Díaz-Galvis J, Schruers K, Berlanga C, Lara-Muñoz C, Griez E. Acute exercise reduces the effects of a 35% CO₂ challenge in patients with panic disorder. *J Affect Disord*. 2007 Aug 27;

Flegal KE, Kishiyama S, Zajdel D, Haas M, Oken BS. Adherence to yoga and exercise interventions in a 6-month clinical trial. *BMC Complement Altern Med*. 2007 Nov 9;7(1):37

Focht, B.; Koltyn, K. Influence of resistance exercise of different intensities on state anxiety and blood pressure. *Med. Sci. Sports Exerc.*, Vol. 31, No. 3, pp. 456-463, 1999.

Fricchione G. Generalized anxiety disorder. *N Engl J Med* 2004; 351:675-82.

Fulk LJ, Stock HS, Lynn A, Marshall J, Wilson MA, Hand GA. Chronic physical exercise reduces anxiety-like behavior in rats. *Int J Sports Med*. 2004 Jan;25(1):78-82.

Galper DI, Trivedi MH, Barlow CE, Dunn AL, Kampert JB. Inverse association between physical inactivity and mental health in men and women. *Med Sci Sports Exerc*. 2006 Jan;38(1):173-8.

Goodwin RD. Association between physical activity and mental disorders among adults in the United States. *Prev Med*. 2003 Jun;36(6):698-703.

Gupta N, Khera S, Vempati RP, Sharma R, Bijlani RL. Effect of yoga based lifestyle intervention on state and trait anxiety. *Indian J Physiol Pharmacol*. 2006 Jan-Mar;50(1):41-7.

Hale, B. S., & Raglin, J. S. (2002). State anxiety responses to acute resistance training and step aerobic exercise across eight weeks of training. *Journal of Sports Medicine and Physical Fitness*, 42, 108– 112.

Kirkwood G, Rampes H, Tuffrey V, Richardson J, Pilkington K. Yoga for anxiety: a systematic review of the research evidence. *Br J Sports Med.* 2005 Dec;39(12):884-91; discussion 891. Review.

Knapen J, Van de Vliet P, Van Coppenolle H, David A, Peuskens J, Pieters G, Knapen K. Comparison of changes in physical self-concept, global self-esteem, depression and anxiety following two different psychomotor therapy programs in nonpsychotic psychiatric inpatients. *Psychother Psychosom.* 2005;74(6):353-61.

Knobf MT, Musanti R, Dorward J. Exercise and quality of life outcomes in patients with cancer. *Semin Oncol Nurs.* 2007 Nov;23(4):285-96.

Larun L, Nordheim LV, Ekeland E, Hagen KB, Heian F. Exercise in prevention and treatment of anxiety and depression among children and young people. *Cochrane Database Syst Rev.* 2006 Jul 19;3:CD004691. Review.

La Forge R. Mind-body fitness: encouraging prospects for primary and secondary prevention. *J Cardiovasc Nurs.* 1997 Apr;11(3):53-65. Review

Lavie CJ, Milani RV. Generalized anxiety disorder. *N Engl J Med.* 2004 Nov 18;351(21):2239; author reply 2239.

Lavie CV, Milani RV. Impact of aging on hostility in coronary patients and effects of cardiac rehabilitation and exercise training in elderly persons. *Am J Geriatr Cardiol.* 2004 May-Jun;13(3):125-30

Lan C, Lai JS, Chen SY, Wong MK. 12-month Tai Chi training in the elderly: its effect on health fitness. *Med Sci Sports Exerc.* 1998 Mar;30(3):345-51.

Larun L, Nordheim LV, Ekeland E, Hagen KB, Heian F. Exercise in prevention and treatment of anxiety and depression among children and young people. *Cochrane Database of Systematic Reviews* 2006, Issue 3. Art. No.: CD004691. DOI: 10.1002/14651858.CD004691.pub2.

LeDoux JE, Gorman JM. A call to action: overcoming anxiety through active coping. *Am J Psychiatry.* 2001 Dec;158(12):1953-5.

Li JX, Hong Y, Chan KM. Tai chi: physiological characteristics and beneficial effects on health. *Br J Sports Med.* 2001 Jun;35(3):148-56. Review.

Martinsen, E.W., Hoffart, A. & Solberg, O. (1989). Aerobic and nonaerobic forms of exercise in the treatment of anxiety disorders. *Stress Medicine*, 5, 115-120.

Mathew SJ, Ho S. Etiology and neurobiology of social anxiety disorder. *J Clin Psychiatry.* 2006;67 Suppl 12:9-13. Review.

Stubbe JH, de Moor MH, Boomsma DI, de Geus EJ. The association between exercise participation and well-being: a co-twin study. *Prev Med.* 2007 Feb;44(2):148-52. Epub 2006 Oct 23.

De Moor MH, Beem AL, Stubbe JH, Boomsma DI, De Geus EJ. Regular exercise, anxiety, depression and personality: a population-based study. *Prev Med.* 2006 Apr;42(4):273-9. Epub 2006 Jan 24.

Morgan, W.P. & O'Connor, P.J. (1987). Exercise and mental health. In R.K. Dishman (Ed.), Exercise Adherence (pp 91-121). Champaign, IL: Human Kinetics Publishers.

Merom D, Phongsavan P, Wagner R, Chey T, Marnane C, Steel Z, Silove D, Bauman A. Promoting walking as an adjunct intervention to group cognitive behavioral therapy for anxiety disorders-A pilot group randomized trial. *J Anxiety Disord.* 2007 Sep 29;

Meyer T, Brooks A. Therapeutic impact of exercise on psychiatric diseases: guidelines for exercise testing and prescription. *Sports Med.* 2000 Oct;30(4):269-79. Review

Motl R., O'Connor P., Dishman R. Effects of cycling exercise on the soleus H-reflex and state anxiety among men with low or high trait anxiety. *Psychophysiology*, 41 (2004), 96–105.

Nouri, S. & Beer, J. (1989). Relations of moderate physical exercise to scores on hostility, aggression, and trait-anxiety. Perceptual and Motor Skills, 68, 1191-1194.

O'Connor, P. J., Raglin, J. S., & Martinsen, E. W. (2000). Physical activity, anxiety and anxiety disorders. *International Journal of Sport Psychology*, 31, 136–155.

O'Connor, P.J., Raglin, J.S. and Martinsen, E.W. (2000) Physical activity, anxiety and anxiety disorders. *International Journal of Sport Psychology* 31, 136-155.

Otto MW, Church TS, Craft LL, Greer TL, Smits JA, Trivedi MH. Exercise for mood and anxiety disorders. *Prim Care Companion J Clin Psychiatry.* 2007;9(4):287-94.

Paluska, S. A. and Schwenk, T.L. (2000) Physical activity and mental health: current concepts. *Sports Medicine* 29, 167-180.

Parente, D. (2000) Influence of aerobic and stretching exercise on anxiety and sensation-seeking mood state. *Perceptual and Motor Skills* 90, 347-348.

Petruzzello, S. J., & Landers, D. M. (1994). State anxiety reduction and exercise: Does hemispheric activation reflect such changes? *Medicine and Science in Sports and Exercise*, 26, 1028–1035.

Petruzzello, S. J., Landers, D. M., Hatfield, B. D., Kubitz, K. A., & Salazar, W. ~1991 A meta-analysis on the anxiety-reducing effects of acute and chronic exercise. *Sports Medicine*, 11, 143–182.

Petruzzello, S. J., Landers, D. M., Hatfield, B. D., Kubitz, K. A., & Salazar, W. (1991). A meta-analysis on the anxiety reducing effects of acute and chronic exercise: Outcomes and mechanisms. *Sports Medicine*, 11, 143–182

Raglin, J. S. and Wilson, M. (1996) State anxiety following 20 minutes of bicycle ergometer exercise at selected intensities. *International Journal of Sports Medicine* 17, 467-471.

Raglin, J.S. (1990) Exercise and mental health: beneficial and detrimental effects. *Sports Medicine* 9, 323- 329.

Sale C, Guppy A, El-Sayed M. Individual differences, exercise and leisure activity in predicting affective well-being in young adults. *Ergonomics*. 2000 Oct;43(10):1689-97

Salmon P. Effects of physical exercise on anxiety, depression, and sensitivity to stress: a unifying theory. *Clin Psychol Rev*. 2001 Feb;21(1):33-61

Scully, D., Kremer J., Meade, M.M., Graham, R. and Dudgeon, K. (1998) Physical exercise and psychological well-being: A critical review. *British Journal of Sports Medicine* 32, 111-120.

Smith C, Hancock H, Blake-Mortimer J, Eckert K. A randomised comparative trial of yoga and relaxation to reduce stress and anxiety. *Complement Ther Med*. 2007 Jun;15(2):77-83.

Snowball, J. and Szabo, A. (1999) Anxiety, affect and exercise: Preliminary evidence lends support to the Distraction Hypothesis. *Journal of Sport Sciences* 17, 67-68.

Soravia LM, Heinrichs M, Aerni A, Maroni C, Schelling G, Ehlert U, Roozendaal B, de Quervain DJ. Glucocorticoids reduce phobic fear in humans. *Proc Natl Acad Sci U S A*. 2006 Apr 4;103(14):5585-90

Spano L The relationship between exercise and anxiety, obsessive compulsiveness, and narcissism. . *Personality and Individual Differences* 30 (2001) 87±93

Steptoe A, Edwards S, Moses J, Mathews A. The effects of exercise training on mood and perceived coping ability in anxious adults from the general population. *J Psychosom Res* 1989;33:537– 47.

Ströhle A, Feller C, Strasburger CJ, Heinz A, Dimeo F. Anxiety modulation by the heart? Aerobic exercise and atrial natriuretic peptide. *Psychoneuroendocrinology*. 2006 Oct;31(9):1127-30.

Ströhle A, Feller C, Onken M, Godemann F, Heinz A, Dimeo F. The acute antipanic activity of aerobic exercise. *Am J Psychiatry*. 2005 Dec;162(12):2376-8

Ströhle A, Holsboer F. Stress responsive neurohormones in depression and anxiety. *Pharmacopsychiatry*. 2003 Nov;36 Suppl 3:S207-14

Sudo, A. (1983). Time course of the changes of catecholamine levels in rat brain during swimming stress. *Brain Research*, 276, 372-374.

Sutoo, D.E. & Akiyama, K. (1996). The mechanism by which exercise modifies brain function. *Physiology Behavior* 60(1), 177-181.

Szabo A., Mesko, A., Caputo, A. and Gill, É. (1998) Examination of exercise-induced feeling states in four modes of exercise. *International Journal of Sport Psychology* 29, 376-390.

Szabo, A. (2003) The acute effects of humor and exercise on mood and anxiety. *Journal of Leisure Research* 35, 152-162.

Tate, A.K. & Petruzello, S.J. (1995). Varying the intensity of acute exercise: implications for changes in affect. *Journal of Sports and Medicine in Physical Fitness* 35(4), 295-302.

Thayer, R. E., Newman, J. R., & McClain, T. M. (1994). Self-regulation of mood: Strategies for changing a bad mood, raising energy, and reducing tension. *Journal of Personality and Social Psychology*, 67, 910–925.

Tieman, J. G., Peacock, L. J., Cureton, K. J., & Dishman, R. K. (2002). State anxiety after acute exercise: Intensity and history. *International Journal of Sport Psychology*, 33, 155–166.

Trejo JL, Llorens-Martín MV, Torres-Alemán I. The effects of exercise on spatial learning and anxiety-like behavior are mediated by an IGF-I-dependent mechanism related to hippocampal neurogenesis. *Mol Cell Neurosci*. 2007 Nov 5

Trivedi MH, Greer TL, Grannemann BD, Church TS, Galper DI, Sunderajan P, Wisniewski SR, Chambliss HO, Jordan AN, Finley C, Carmody TI. TREAD: Treatment with Exercise Augmentation for Depression: study rationale and design. *Clin Trials*. 2006;3(3):291-305.

Wilson, V.E., Berger, B.G. and Bird, E.I. (1981) Effects of running and of an exercise class on anxiety. *Perceptual and Motor Skills* 53, 472-474.

Wininger SR. Improvement of affect following exercise: methodological artifact or real finding? *Anxiety Stress Coping*. 2007 Mar;20(1):93-102.