

Chapter 5 Depression

Adlard, P.A., Perreau, V.M., & Cotman, C.W. (2005). The exercise-induced expression of BDNF within the hippocampus varies across life-span. *Neurobiology of Aging*, 26, 511-520.

Adlard, P.A., Perreau, V.M., Engesser-Cesar, C., & Cotman, C.W. (2004). The time course of induction of brain-derived neurotrophic factor mRNA and protein in the rat hippocampus following voluntary exercise. *Neuroscience Letters*, 363, 43-48.

Akama KT, McEwen BS. Gene therapy to bet on: protecting neurons from stress hormones. *Trends Pharmacol Sci*. 2005 Apr;26(4):169-72. Review.

Ames, E. The development of Romanian orphanage children adopted to Canada (Final Report to the National Welfare Grants Program; Human Resources Development. Canada) Burnaby, Canada; Simon Fraser University (1997)

Babyak, M. A., Blumenthal, J. A., Herman, S., Khatri, P., Doraiswamy, P. M., Moore, K. A., Craighead, W. E., et al. (2000). Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosomatic Medicine*, 62, 633-638.

Barbour KA, Blumenthal JA. Exercise training and depression in older adults. *Neurobiol Aging*. 2005 Dec;26 Suppl 1:119-23. Epub 2005 Oct 11. Review.

Barbour KA, Edenfield TM, Blumenthal JA. Exercise as a Treatment for Depression and Other Psychiatric Disorders: A REVIEW. *J Cardiopulm Rehabil Prev*. 2007 Nov-Dec;27(6):359-67.

Berchtold NC, Chinn G, Chou M, Kesslak JP, Cotman CW. Exercise primes a molecular memory for brain-derived neurotrophic factor protein induction in the rat hippocampus. *Neuroscience*. 2005;133(3):853-61.

Berger, B.G. and Motl, R.W. (2000) Exercise and mood: A selective review and synthesis of research employing the Profile of Mood States. *Journal of Applied Sport Psychology* 12, 69-92.

Berger, B.G. and Owen, D.R. (1992) Mood alteration with yoga and swimming; aerobic exercise may not be necessary. *Perceptual and Motor Skills* 75,1331-1343.

Birmaher B, Ryan ND, Williamson DE, Brent DA, Kaufman J, Dahl RE, Perel J, Nelson B. Childhood and adolescent depression: A review of the past 10 years. Part I. *J Am Acad Child Adolesc Psychiatry* 1996;35: 1427-39.

Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, Waugh R, Napolitano MA, Forman LM, Appelbaum M, Doraiswamy PM, Krishnan KR. Effects of exercise training on older patients with major depression. *Arch Intern Med* 1999;159:2349-56.

Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, Herman S, Craighead WE, Brosse AL, Waugh R, Hinderliter A, Sherwood A. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*. 2007 Sep-Oct;69(7):587-96. Epub 2007 Sep 10.

Brené S, Bjørnebekk A, Aberg E, Mathé AA, Olson L, Werme M. Running is rewarding and antidepressive. *Physiol Behav*. 2007 Sep 10;92(1-2):136-40. Epub 2007 May 21.

Brown JD, Lawton M. Stress and well-being in adolescence: The moderating role of physical exercise. *J Hum Stress* 1986;12:125–31.

Brown SW, Welsh MC, Labbe EE, Vitulli WF, Kulkarni P. Aerobic exercise in the psychological treatment of adolescents. *Percept Mot Skills* 1982;74:555–60.

Byrne, A., & Byrne, D. G. (1993). The effect of exercise on depression, anxiety and other mood states. *Journal of Psychosomatic Research*, 37(6), 565-574.

Camacho, T. C., Roberts, R. E., Lazarus, N. B., Kaplan, G. A., & Cohen, R. D. (1991). Physical activity and depression: Evidence from the Alameda County Study. *American Journal of Epidemiology*, 134, 220-231.

Campbell S, Macqueen G. The role of the hippocampus in the pathophysiology of major depression. *J Psychiatry Neurosci*. 2004 Nov;29(6):417-26. Review.

Chen MJ, Russo-Neustadt AA. Running exercise- and antidepressant-induced increases in growth and survival-associated signaling molecules are IGF-dependent. *Growth Factors*. 2007 Apr;25(2):118-31.

Cooper-Patrick, L., Ford, D. E., Mead, L. A., Chang, P. P., & Klag, M. J. (1997). Exercise and depression in midlife: A prospective study. *American Journal of Public Health*, 87(4), 670-673.

Cotman CW, Berchtold NC, Christie LA. Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends Neurosci*. 2007 Sep;30(9):464-72. Epub 2007 Aug 31.

Cross-National Collaborative Group. The changing rate of major depression. *JAMA* 1992;268:3098–105.

depression among older adults. *Alternative Therapies in Health and Medicine*, 4(1),

DiLorenzo TM, Bargman EP, Stucky-Ropp R, Brassington GS, Frensch PA, LaFontaine T. Long-term effects of aerobic exercise on psychological outcomes. *Prev Med* 1999;28:75– 85.

Ding Q, Vaynman S, Akhavan M, Ying Z, Gomez-Pinilla F. Insulin-like growth factor I interfaces with brain-derived neurotrophic factor-mediated synaptic plasticity to modulate aspects of exercise-induced cognitive function. *Neuroscience*. 2006 Jul 7;140(3):823-33. Epub 2006 May

Doyne EJ, Ossip-Klein DJ, Bowman ED, Osborn KM, McDougall- Wilson IB, Neimeyer RA. Running versus weight lifting in the treatment of depression. *J Consult Clin Psychol* 1987;55:748– 54.

Drevets WC. Neuroimaging studies of mood disorders. *Biol Psychiatry*. 2000 Oct 15;48(8):813-29

Duman R, Heninger GR, Nestler EJ (1997): A molecular and cellular theory of depression. *Arch Gen Psychiatry* 54:597–606.

Duman RS, Malberg J, Nakagawa S, D'Sa C (2000): Neuronal plasticity and survival in mood disorders. *Biol Psychiatry* 46:1181–1191.

Duman RS, Malberg J, Nakagawa S, D'Sa C. Neuronal plasticity and survival in mood disorders. *Biol Psychiatry*. 2000 Oct 15;48(8):732-9. Beneficial effects of exercise and its molecular mechanisms on depression in rats. *Behav Brain Res*. 2006 Mar 15;168(1):47-55.

Duman RS, Malberg J, Thome J. Neural plasticity to stress and antidepressant treatment. *Biol Psychiatry*. 1999 Nov 1;46(9):1181-91. Review.

Duman RS. Neurotrophic factors and regulation of mood: role of exercise, diet and metabolism. *Neurobiol Aging*. 2005 Dec;26 Suppl 1:88-93. Epub 2005 Oct 13. Review.

Duman RS. Depression: a case of neuronal life and death? *Biol Psychiatry*. 2004 Aug 1;56(3):140-5. Review.

Duman RS. Neurotrophic factors and regulation of mood: role of exercise, diet and metabolism. *Neurobiol Aging*. 2005 Dec;26 Suppl 1:88-93. Epub 2005 Oct 13. Review.\

Duman RS. Depression: a case of neuronal life and death? *Biol Psychiatry*. 2004 Aug 1;56(3):140-5. Review.

Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. The DOSE study: a clinical trial to examine efficacy and dose response of exercise as treatment for depression. *Control Clin Trials*. 2002 Oct;23(5):584-603.

Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc* 2001;33:S587–97.

Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc*. 2001 Jun;33(6 Suppl):S587-97; discussion 609-10.

Dunn, A. L., Trivedi, M. H., & O'Neil, H. (2001). Physical activity dose-response effects on outcomes of depression and anxiety. *Medicine and Science in Sports and Exercise*, 33(Suppl.), S587-S597.

Dunn, A.L. and Dishman, R.K. (1991) Exercise and the neurobiology of depression. *Exercise and Sport Sciences Reviews* 19, 41-98.

Ekkekakis, P. and Petruzzello, S.J. (1999) Acute aerobic exercise and affect: Current status, problems and prospects regarding dose-response. *Sports Medicine* 28, 337-374.

Ernst C, Olson AK, Pineda JP, Lam RW, Christie BR. Antidepressant effects of exercise: evidence for an adult-neurogenesis hypothesis?

Ernst C, Olson AK, Pineda JP, Lam RW, Christie BR. Antidepressant effects of exercise: evidence for an adult-neurogenesis hypothesis? *J Psychiatry Neurosci*. 2006 Mar;31(2):84-92. Review.

Exercise referral and the treatment of mild or moderate depression Information for GPs and healthcare practitioners. 2006
<http://www.mentalhealth.org.uk/EasysiteWeb/getresource.axd?AssetID=39360&type=Full&servicetype=Attachment>.

Fabel, K., Fabel, K, Tam, B., Kaufer, D., Baiker, A., Simmons, N., Kuo, C.J., Palmer, T.D. (2003). VEGF is necessary for exercise-induced adult hippocampal neurogenesis. *European Journal of Neuroscience*, 18, 2803-2812.

Farrell, P.A., Gates, W.K., Maksud, M.G. and Morgan, W.P. (1982). Increase in plasma betaendorphin beta-lipotropin immunoreactivity after treadmill running in humans. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology* 52, 1245-1249.

Farrell, P.A., Gustafson, A.B., Morgan, W.P. and Pert, C.B. (1987) Enkephalins, catecholamines, and psychological mood alterations: effects of prolonged exercise. *Medicine and Science in Sports and Exercise*, 19, 347-353.

Field T, Diego M, Sanders CE. Exercise is positively related to adolescents' relationships and academics. *Adolescence* 2001;36:105-10.

Fleming JE, Offord DR. Epidemiology of childhood depressive disorders: A critical review. *J Am Acad Child Adolesc Psychiatry* 1990;29:571-80.

Fremont J, Craighead LW. Aerobic exercise and cognitive therapy in the treatment of dysphoric moods. *Cognitive Ther Res* 1987;11:241-51.

Frodl T, Meisenzahl EM, Zetsche T, Born C, Groll C, Jäger M, et al (2002): Hippocampal changes in patients with a first episode of major depression. *Am J Psychiatry* 159:1112-1118.

Garcia C, Chen MJ, Garza AA, Cotman CW, Russo-Neustadt A. The influence of specific noradrenergic and serotonergic lesions on the expression of hippocampal brain-derived neurotrophic factor transcripts following voluntary physical activity. *Neuroscience*. 2003;119(3):721-32.

Garza AA, Ha TG, Garcia C, Chen MJ, Russo-Neustadt AA. Exercise, antidepressant treatment, and BDNF mRNA expression in the aging brain. *Pharmacol Biochem Behav*. 2004 Feb;77(2):209-20.

Garza AA, Ha TG, Garcia C, Chen MJ, Russo-Neustadt AA. Exercise, antidepressant treatment, and BDNF mRNA expression in the aging brain. *Pharmacol Biochem Behav*. 2004 Feb;77(2):209-20.

Greist, J. H., Klein, M. H., Eischens, R. R., Gurman, A. S., & Morgan, W. J. (1979). Running as treatment for depression. *Comprehensive Psychiatry*, 29, 41-54.

Gunnar, M. et al *Development and Psychopathology* 13: 611 – 628 (2001)

Harland J, White M, Drinkwater C, Chinn D, Farr L, Howel D. The Newcastle exercise project: a randomized controlled trial of methods to promote physical activity in primary care.

Herman S, Blumenthal JA, Babyak M, Khatri P, Craighead WE, Krishnan KR, Doraiswamy PM. Exercise therapy for depression in middle-aged and older adults: predictors of early dropout and treatment failure. *Health Psychol*. 2002 Nov;21(6):553-63.

Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur J Neurosci*. 2004 Nov;20(10):2580-90

Ivy AS, Rodriguez FG, Garcia C, Chen MJ, Russo-Neustadt AA. Noradrenergic and serotonergic blockade inhibits BDNF mRNA activation following exercise and antidepressant. *Pharmacol Biochem Behav*. 2003 Apr;75(1):81-8.

J Psychiatry Neurosci. 2006 Mar;31(2):84-92.

Jacobs BL, Praag H, Gage FH (2000): Adult brain neurogenesis and psychiatry: A novel theory of depression. *Mol Psychiatry* 5:262–269.

Kelder SH, Perry CL, Klepp KI, Lytle LL. Longitudinal tracking of adolescent smoking, physical activity, and food choice behaviors. *Am J Public Health* 1994;84:1121–6.

Kempermann G (2002): Regulation of adult hippocampal neurogenesis—implications for novel theories of major depression. *Bipolar Disord* 4:17–33.

Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen H, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994;51:8–18.

Kessler RC, Walters EE. Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Co-morbidity Survey. *Depress Anxiety* 1998;7:3–14.

Kirkcaldy BD, Shephard RJ, Siefen RG. The relationship between physical activity and self-image and problem behaviour among adolescents. *Soc Psychiatry Psychiatr Epidemiol* 2002;37:544–50.

Kovacs M. Psychiatric disorders in youths with IDDM: Rates and risk factors. *Diabetes Care* 1997;20:36–44.

Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: Systematic review and meta-regression analysis of randomised controlled trials. *Brit Med J* 2001;322:1–8.

Lee AL, Ogle WO, Sapolsky RM. Stress and depression: possible links to neuron death in the hippocampus. *Bipolar Disord.* 2002 Apr;4(2):117-28. Review.

Lewinsohn PM, Rohde P, Seeley JR. Major depressive disorder in older adolescents: Prevalence, risk factors, and clinical implications. *Clin Psych Rev* 1998;18:765–94.

Madsen TM, Treschow A, Gengzon J, Bolwig TG, Lindvall O, Tingström A (2000): Increased neurogenesis in a model of electroconvulsive therapy. *Biol Psychiatry* 47:1043–1049.

Manji HK, Duman RS. Impairments of neuroplasticity and cellular resilience in severe mood disorders: implications for the development of novel therapeutics. *Psychopharmacol Bull.* 2001 Spring;35(2):5-49. Review.

Manji HK, Quiroz JA, Sporn J, Payne JL, Denicoff K, A Gray N, Zarate CA Jr, Charney DS. Enhancing neuronal plasticity and cellular resilience to develop novel, improved therapeutics for difficult-to-treat depression. *Biol Psychiatry.* 2003 Apr 15;53(8):707-42. Review

Martinsen, E. W. ~1990!. Benefits of exercise for the treatment of depression.

Martinsen, E. W., Hoffart, A., & Solberg, O. (1989). Comparing aerobic and nonaerobic forms of exercise in the treatment of clinical depression: A randomized trial. *Comprehensive Psychiatry*, 30(4), 324-331.

Martinsen, E.W., Medhus, A., & Sandvik, L. (1985). Effects of aerobic exercise on depression: A controlled study. *British Medical Journal*, 291, 109.

Martinsen, E.W., Sandvik, L., & Kolbjornsrud, O. B. (1989). Aerobic exercise in the treatment of nonpsychotic mental disorders. *Norwegian Journal of Psychiatry*, 43, 411-415.

McDermott RJ, Hawkins WE, Marty PJ, Littlefield EA, Murray S, Williams TK. Health behavior correlates of depression in a sample of high school students. *J School Health* 1990;60:414-7.

McEwen BS, Magarinos AM, Reagan LP. Structural plasticity and tianeptine: cellular and molecular targets. *Eur Psychiatry*. 2002 Jul;17 Suppl 3:318-30. Review.

McEwen BS, Olie JP. Neurobiology of mood, anxiety, and emotions as revealed by studies of a unique antidepressant: tianeptine.

McEwen BS. Glucocorticoids, depression, and mood disorders: structural remodeling in the brain. *Metabolism*. 2005 May;54(5 Suppl 1):20-3. Review.

McEwen BS. Glucocorticoids, depression, and mood disorders: structural remodeling in the brain. *Metabolism*. 2005 May;54(5 Suppl 1):20-3. Review.

McKittrick CR, Magariños AM, Blanchard DC, Blanchard RJ, McEwen BS, Sakai RR. Chronic social stress reduces dendritic arbors in CA3 of hippocampus and decreases binding to serotonin transporter sites. *Synapse*. 2000 May;36(2):85-94.

Mechanic D, Hansell S. Adolescent competence, psychological wellbeing, and self-assessed physical health. *J Health Soc Behav* 1987;28: 364-74.

Mitra R, Sundlass K, Parker KJ, Schatzberg AF, Lyons DM. Social stress-related behavior affects hippocampal cell proliferation in mice. *Physiol Behav*. 2006 Sep 30;89(2):123-7

Mol Psychiatry. 2005 Jun;10(6):525-37. Review.

Molteni R, Wu A, Vaynman S, Ying Z, Barnard RJ, Gómez-Pinilla Exercise reverses the harmful effects of consumption of a high-fat diet on synaptic and behavioral plasticity associated to the action of brain-derived neurotrophic factor. *Neuroscience*. 2004;123(2):429-40.

Moore KA, Blumenthal JA. Exercise training as an alternative treatment for depression among older adults. *Altern Ther Health Med* 1998;4: 48-56.

Moore, K. A., & Blumenthal, J. A. (1998). Exercise training as an alternative treatment for

Morgan, K., & Bath, P. A. (1998). Customary physical activity and psychological wellbeing. *Age and Ageing*, 27, 35-40.

Morgan, W.P. (1985) Affective beneficence of vigorous physical activity. *Medicine and Science in Sports and Exercise* 17, 94-100.

Morgan, W.P. and O'Connor, P.J. (1988). Exercise and mental health. In: Exercise Adherence: Its impact on Public Health. Ed: Dishman R.K. Champaign, IL: Human Kinetics. 91-121.

Morgan, W. P. (1969). Physical fitness and emotional health. *American Corrective Therapy Journal*, 23, 124-127.

Morgan, W. P. (1994). Physical activity, fitness and depression. In T. Stephens (Ed.), *Physical activity, fitness and health* (pp. 851-867). Champaign, IL: Human Kinetics.

Morgan, W. P., Roberts, J. A., Brand, F. R., & Feinerman, A. D. (1970). Psychologic effect of chronic physical activity. *Medicine and Science in Sports and Exercise*, 2, 213-217.

Murray CJL, Lopez AD. Evidence-based health policy – lessons from the Global Burden of Disease Study. *Science* 1996;274:740–3.

Murray NG, Kelder SH, Meyers MC. Type of physical activity and level of depression in suburban 9th and 10th grade students. *Med Sci Sports Exerc* 1998;30:S117.

Nesse RM. Is depression an adaptation? *Arch Gen Psychiatry* 2000;57:14-20.

Neumeister A, Wood S, Bonne O, Nugent AC, Luckenbaugh DA, Young T, Bain EE, Charney DS, Drevets WC. Reduced hippocampal volume in unmedicated, remitted patients with major depression versus control subjects. *Biol Psychiatry*. 2005 Apr 15;57(8):935-7

Niculescu AB. Genomic studies of mood disorders -- the brain as a muscle? *Genome Biol*. 2005;6(4):215. Epub 2005 Mar 24. Review.

Olson AK, Eadie BD, Ernst C, Christie BR. Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. *Hippocampus*. 2006;16(3):250-60. Review.

Olson AK, Eadie BD, Ernst C, Christie BR. Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. *Hippocampus*. 2006;16(3):250-60. Review.

Page RM, Tucker LA. Psychosocial discomfort and exercise frequency: An epidemiological study of adolescents. *Adolescence* 1994;29:183–91.

Pate RR, Heath GW, Dowda M, Trost SG. Associations between physical activity and other health behaviors in a representative sample of US adolescents. *Am J Public Health* 1996;86:1577–81.

Rosenzweig, M. R. & Bennett, E. L. (1996). Psychobiology of plasticity: effects of training and experience on brain and behavior. *Behavioral Brain Research*, 78, 57-65.

- Rushton JL, Forcier M, Schectman RM. Epidemiology of depressive symptoms in the National Longitudinal Study of Adolescent Health. *J Am Acad Child Adolesc Psychiatry* 2002;41:199–205.
- Russo-Neustadt A, Beard RC, Cotman CW. Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropsychopharmacology*. 1999 Nov;21(5):679-82.
- Russo-Neustadt A, Ha T, Ramirez R, Kessler JP. Physical activity-antidepressant treatment combination: impact on brain-derived neurotrophic factor and behavior in an animal model. *Behav Brain Res*. 2001 Apr 8;120(1):87-95.
- Russo-Neustadt A. Brain-derived neurotrophic factor, behavior, and new directions for the treatment of mental disorders. *Semin Clin Neuropsychiatry*. 2003 Apr;8(2):109-18. Review.
- Russo-Neustadt AA, Alejandre H, Garcia C, Ivy AS, Chen MJ. Hippocampal brain-derived neurotrophic factor expression following treatment with reboxetine, citalopram, and physical exercise. *Neuropsychopharmacology*. 2004 Dec;29(12):2189-99.
- Russo-Neustadt AA, Beard RC, Huang YM, Cotman C. Physical activity and antidepressant treatment potentiate the expression of specific brain-derived neurotrophic factor transcripts in the rat hippocampus. *Neuroscience*. 2000;101(2):305-12.
- Russo-Neustadt AA, Chen MJ. Brain-derived neurotrophic factor and antidepressant activity. *Curr Pharm Des*. 2005;11(12):1495-510. Review
- Russo-Neustadt AA, Chen MJ. Brain-derived neurotrophic factor and antidepressant activity. *Curr Pharm Des*. 2005;11(12):1495-510. Review. 48-56
- Russo-Neustadt, A., Ryan, C.B. & Cotman, C.W. (1999). Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropsychopharmacology*, 21, 679-682.
- Ryan ND, Puig-Antich J, Ambrosini P, Rabinovich H, Robinson D, Nelson B, Iyengar S, Twomey J. The clinical picture of major depression in children and adolescents. *Arch Gen Psychiatry* 1987;44:854–61.
- Sanders CE, Field TM, Diego M, Kaplan M. Moderate involvement in sports is related to lower depression levels among adolescents. *Adolescence* 2000;35:793–7.
- Sapolsky RM. Chickens, eggs and hippocampal atrophy. *Nat Neurosci*. 2002 Nov;5(11):1111-3.
- Sapolsky RM. Is impaired neurogenesis relevant to the affective symptoms of depression? *Biol Psychiatry*. 2004 Aug 1;56(3):137-9

Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature*. 2004 May 13;429(6988):184-7

Schmitz KH, Lytle LA, Phillips GA, Murray DM, Birnbaum AS, Kubik MY. Psychosocial correlates of physical activity and sedentary leisure habits in young adolescents: The Teens Eating for Energy and Nutrition study. *Prev Med* 2002;34:266–78.

Shaffer, D., Gould, M. S., Fisher, P., Trautman, P., Moreau, D., Kleinman, M., Flory M. Psychiatric diagnosis in child and adolescent suicide. *Arch Gen Psychiatry* 1996;53:339–48.

Sharp T. Neuroscience. A new molecule to brighten the mood. *Science*. 2006 Jan 6;311(5757):45-6.

Singh NA, Clements KM, Singh MA. The efficacy of exercise as a long-term antidepressant in elderly subjects: a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci* 2001;56:M497–504.

Sports Medicine, 9, 380–389.

Stein DJ, Collins M, Daniels W, Noakes TD, Zigmond M. Mind and muscle: the cognitive-affective neuroscience of exercise. *CNS Spectr*. 2007 Jan;12(1):19-22.

Stephens, T. (1988). Physical activity and mental health in the United States and Canada: Evidence from four population surveys. *Preventive Medicine*, 17, 35-47.

Stephens, T., & Craig, C. L. (1990). *The well-being of Canadians: Highlights of the 1988 Cambell's Survey*. Ottawa: Canadian Fitness and Lifestyle Research Institute.

Steptoe A, Butler N. Sports participation and emotional well-being in adolescents. *Lancet* 1996;347:1789–92.

Stoll, O. (1997) Endorphine, Laufsucht und runner's high. Aufstieg und niedergang eines mythos (Endogenous opiates, "runner's high" and "Exercise Addiction" - The rise and decline of a myth). *Leipziger Sportwissenschaftliche Beitrage* 28, 102-121.

Stranahan AM, Khalil D, Gould E. Social isolation delays the positive effects of running on adult neurogenesis. *Nat Neurosci*. 2006 Apr;9(4):526-33. Epub 2006 Mar 12.

Strawbridge WJ, Deleger S, Roberts RE, Kaplan GA. Physical Activity Reduces the Risk of Subsequent Depression for Older Adults *Am J Epidemiol* 2002;156:328–34.

Szabo A., Mesko, A., Caputo, A. and Gill, É. (1998) Examination of exercise-induced feeling states in four modes of exercise. *International Journal of Sport Psychology* 29, 376-390.

Szabo, A. (2003) The acute effects of humor and exercise on mood and anxiety. *Journal of Leisure Research* 35, 152-162.

Szabo A. Acute psychological effects of exercise performed. at self-selected workloads: Implications for theory and practice.. *Journal of Sports Science and Medicine* (2003) 2, 77-87

Vaidya VA, Siuciak JA, Du F, Duman RS. Hippocampal mossy fiber sprouting induced by chronic electroconvulsive seizures. *Neuroscience*. 1999 Mar;89(1):157-66.

Van Praag H, Shubert T, Zhao C, Gage FH. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci*. 2005 Sep 21;25(38):8680-5

Van Praag H, Shubert T, Zhao C, Gage FH. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci*. 2005 Sep 21;25(38):8680-5.

Vaynman S, Gomez-Pinilla F. License to run: exercise impacts functional plasticity in the intact and injured central nervous system by using neurotrophins. *Neurorehabil Neural Repair*. 2005 Dec;19(4):283-95. Review.

Vaynman S, Gomez-Pinilla F. Revenge of the "sit": how lifestyle impacts neuronal and cognitive health through molecular systems that interface energy metabolism with neuronal plasticity. *J Neurosci Res*. 2006 Sep;84(4):699-715. Review.

Vaynman S, Ying Z, Gomez-Pinilla F. Interplay between brain-derived neurotrophic factor and signal transduction modulators in the regulation of the effects of exercise on synaptic-plasticity. *Neuroscience*. 2003;122(3):647-57

Vaynman S, Ying Z, Wu A, Gomez-Pinilla F. Coupling energy metabolism with a mechanism to support brain-derived neurotrophic factor-mediated synaptic plasticity. *Neuroscience*. 2006;139(4):1221-34. Epub 2006 Mar 31. PMID: 16580138

Vaynman SS, Ying Z, Yin D, Gomez-Pinilla F. Exercise differentially regulates synaptic proteins associated to the function of BDNF. *Brain Res*. 2006 Jan 27;1070(1):124-30. Epub 2006 Jan 18. Vaynman S, Ying Z, Gomez-Pinilla F. Abstract

Vollmayr B, Simonis C, Weber S, Gass P, Henn FA (2003): Reduced cell proliferation in the dentate gyrus is not correlated with the development of learned helplessness. *Biol Psychiatry* 54:1035–1040.

Warner-Schmidt JL, Duman RS. Hippocampal neurogenesis: opposing effects of stress and antidepressant treatment. *Hippocampus*. 2006;16(3):239-49. Review.

Warner-Schmidt JL, Duman RS. VEGF is an essential mediator of the neurogenic and behavioral actions of antidepressants. *Proc Natl Acad Sci U S A*. 2007 Mar 13;104(11):4647-52. Epub 2007 Mar 5.

Wells VE, Deykin EY, Klerman GL. Risk factors for depression in adolescence. *Psychiatric Devel* 1985;3:83–108.

Zarate CA Jr, Du J, Quiroz J, Gray NA, Denicoff KD, Singh J, Charney DS, Manji HK. Regulation of cellular plasticity cascades in the pathophysiology and treatment of mood disorders: role of the glutamatergic system. *Ann N Y Acad Sci.* 2003 Nov;1003:273-91. Review.

Zarate CA Jr, Singh J, Manji HK. Cellular plasticity cascades: targets for the development of novel therapeutics for bipolar disorder. *Biol Psychiatry.* 2006 Jun 1;59(11):1006-20. Epub 2006 Feb 17.

Zheng H, Liu Y, Li W, Yang B, Chen D, Wang X, Jiang Z, Wang H, Wang Z, Cornelisson G, Halberg F.