

Chapter 8 Women

Asikainen TM, Kukkonen-Harjula K, Miilunpalo S. Exercise for health for early postmenopausal women: a systematic review of randomised controlled trials. *Sports Med.* 2004;34(11):753-78. Review.

Bergmann A, Zygmunt M, Clapp JF 3rd. Running throughout pregnancy: effect on placental villous vascular volume and cell proliferation. *Placenta.* 2004 Sep-Oct;25(8-9):694-8.

Bick-Sander A, Steiner B, Wolf SA, Babu H, Kempermann G. Running in pregnancy transiently increases postnatal hippocampal neurogenesis in the offspring. *Proc Natl Acad Sci U S A.* 2006 Mar 7;103(10):3852-7.

Berchtold NC, Kesslak JP, Pike CJ, Adlard PA, Cotman CW. Estrogen and exercise interact to regulate brain-derived neurotrophic factor mRNA and protein expression in the hippocampus. *Eur J Neurosci.* 2001 Dec;14(12):1992-2002.

Brockie J. Exercise for women in the early postmenopausal years. *J Br Menopause Soc.* 2006 Sep;12(3):126-7.

Bungum TJ, Peaslee DL, Jackson AW, Perez MA. Exercise during pregnancy and type of delivery in nulliparae. *J Obstet Gynecol Neonatal Nurs* 2000;29:258-64.

Busch AJ, Barber KA, Overend TJ, Peloso PM, Schachter CL. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev.* 2007 Oct 17;(4):CD003786. Review.

Christie BR, Swann SE, Fox CJ, Froc D, Lieblich SE, Redila V, Webber A. Voluntary exercise rescues deficits in spatial memory and long-term potentiation in prenatal ethanol-exposed male rats. *Eur J Neurosci.* 2005 Mar;21(6):1719-26.

Clapp JF 3rd, Kim H, Burciu B, Lopez B. Beginning regular exercise in early pregnancy: effect on fetoplacental growth. *Am J Obstet Gynecol.* 2000 Dec;183(6):1484-8

Clapp JF 3rd, Lopez B, Harcar-Sevcik R. Neonatal behavioral profile of the offspring of women who continued to exercise regularly throughout pregnancy. *Am J Obstet Gynecol.* 1999 Jan;180(1 Pt 1):91-4.

Clapp JF 3rd, Simonian S, Lopez B, Appleby-Wineberg S, Harcar-Sevcik R. The one-year morphometric and neurodevelopmental outcome of the offspring of women who continued to exercise regularly throughout pregnancy. *Am J Obstet Gynecol.* 1998 Mar;178(3):594-9.

Clapp JF 3rd. A clinical approach to exercise during pregnancy. *Clin Sports Med.* 1994 Apr;13(2):443-58. Review.

Clapp JF 3rd. Exercise during pregnancy. A clinical update.

Clapp JF 3rd. Morphometric and neurodevelopmental outcome at age five years of the offspring of women who continued to exercise regularly throughout pregnancy. *J Pediatr.* 1996 Dec;129(6):856-63.

Clapp JF 3rd. The effect of continuing regular endurance exercise on the physiologic adaptations to pregnancy and pregnancy outcome. *Am J Sports Med.* 1996;24(6 Suppl):S28-9.

Clapp JF. *Exercising Through Your Pregnancy.* Addicus Books; 1 edition (January 1, 2002).

Clapp JF. Influence of endurance exercise and diet on human placental development and fetal growth. *Placenta.* 2006 Jun-Jul;27(6-7):527-34. Epub 2005 Sep 13. Review.

Clin Sports Med. 2000 Apr;19(2):273-86. Review.

Cook, D, Nagelkirk P., et al. Exercise and Cognitive Performance in Chronic Fatigue Syndrome. *Med. Sci. Sports Exerc.*, Vol. 37, No. 9, pp. 1460–1467, 2005.

Coe CL, Kramer M, Czeh B, Gould E, Reeves AJ, Kirschbaum C, Fuchs E. Prenatal stress diminishes neurogenesis in the dentate gyrus of juvenile rhesus monkeys. *Biol Psychiatry.* 2003 Nov 15;54(10):1025-34.

Crews FT, Nixon K, Wilkie ME. Exercise reverses ethanol inhibition of neural stem cell proliferation. *Alcohol.* 2004 May;33(1):63-71.

Dennerstein L, Lehert P, Burger H, Dudley E. Mood and the menopausal transition. *J Nerv Ment Dis.* 1999 Nov;187(11):685-91.

Elavsky S, McAuley E. Physical activity, symptoms, esteem, and life satisfaction during menopause. *Maturitas.* 2005 Nov-Dec;52(3-4):374-85.

Ezmerli NM. Exercise in pregnancy. *Prim Care Update Ob Gyns* 2000;7:260–5.

Erickson, KI, Colcombe, SJ, Elavsky, S., McAuley, E, Korol, DL, Scalf, PE, Kramer, AF. (2007). Interactive effects of hormone treatment on brain health in postmenopausal women. *Neurobiology of Aging*, 28(2): 179-185.

Erickson, KI, Pruis, TA, Debrey, SM, Bohacek, J., Korol, DL. (2006). Estrogen and exercise interact to up-regulate BDNF levels in the hippocampus but not striatum of middle-aged Brown-Norway rats. Program No. 266.17. Society for Neuroscience Abstracts, 2006.

Etnier JL, Sibley BL. Physical activity and hormone-replacement therapy: interactive effects on cognition? *J Aging Phys Act.* 2004 Oct;12(4):554-67.

Friedenreich, C.M. Physical activity and breast cancer risk: The effect of menopausal status.

Exerc. Sport Sci. Rev., Vol. 32, No. 4, pp. 180–184, 2004.

Fugh-Berman A, Kronenberg F. Complementary and alternative medicine (CAM) in reproductive-age women: a review of randomized controlled trials. *Reprod Toxicol*. 2003 Mar-Apr;17(2):137-52.

Greisen MH, Altar CA, Bolwig TG, Whitehead R, Wörtwein G. Increased adult hippocampal brain-derived neurotrophic factor and normal levels of neurogenesis in maternal separation rats. *J Neurosci Res*. 2005 Mar 15;79(6):772-8.

Halbreich U, Kahn LS (2001) Role of estrogen in the aetiology and treatment of mood disorders. *CNS Drugs* 15(10):797–817

Hartmann S, Bung P. Physical exercise during pregnancy— physiological considerations and recommendations. *J Perinat Med* 1999;27:204–15.

Hartmann S, Bung P, Schlebusch H, Hollmann W. The analgesic effect of exercise during labor. *Z Geburtshilfe Neonatol* 2005 Aug;209(4):144-50.

Heffernan AE. Exercise and pregnancy in primary care. *Nurse Pract* 2000;49:53–6.

Hegaard HK, Pedersen BK, Nielsen BB, Damm P. Leisure time physical activity during pregnancy and impact on gestational diabetes mellitus, pre-eclampsia, preterm delivery and birth weight: a review. *Acta Obstet Gynecol Scand*. 2007;86(11):1290-6.

Katz VL. Water exercise in pregnancy. *Semin Perinatol* 1996;20: 285–91.\

Kemmler W, Engelke K, von Stengel S, Weineck J, Lauber D, Kalender WA. Long-term four-year exercise has a positive effect on menopausal risk factors: the Erlangen Fitness Osteoporosis Prevention Study. *J Strength Cond Res*. 2007 Feb;21(1):232-9.

Kemmler W, Lauber D, Weineck J, Hensen J, Kalender W, Engelke K. Benefits of 2 years of intense exercise on bone density, physical fitness, and blood lipids in early postmenopausal osteopenic women: results of the Erlangen Fitness Osteoporosis Prevention Study (EFOPS). *Arch Intern Med*. 2004 May 24;164(10):1084-91.

Keizer HA, Kuipers H et al (1987) Effect of a 3-month endurance training program on metabolic and multiple hormonal responses to exercise. *Int J Sports Med* 8(Suppl 3)(12):154–160.

Koltyn, K.F. and Schultes, S.S. (1997) Psychological effects of an aerobic exercise session and a rest session following pregnancy. *Journal of Sports Medicine and Physical Fitness* 37, 287-291.

Korol, DL & Pruis, TA (2004). Estrogen and exercise modulate learning strategy in middle-aged female rats. Program No. 770.7. Society for Neuroscience Abstracts, 2004.

Korol, DL & Pruis, TA (2004). Estrogen and exercise modulate learning strategy in middle-aged female rats. Program No. 770.7. Society for Neuroscience Abstracts, 2004.

Kritz-Silverstein D, Wingard DL, Garland FC. The association of behavior and lifestyle factors with menstrual symptoms. *J Womens Health Gend Based Med.* 1999 Nov;8(9):1185-93.

Kroll R, Rapkin AJ. Treatment of premenstrual disorders. *J Reprod Med.* 2006 Apr;51(4 Suppl):359-70

Lee HH, Kim H, Lee JW, Kim YS, Yang HY, Chang HK, Lee TH, Shin MC, Lee MH, Shin MS, Park S, Baek S, Kim CJ. Maternal swimming during pregnancy enhances short-term memory and neurogenesis in the hippocampus of rat pups. *Brain Dev.* 2006 Apr;28(3):147-54. Epub 2005 Dec 20.

Lind E, Joens-Matre RR, Ekkekakis P. What intensity of physical activity do previously sedentary middle-aged women select? Evidence of a coherent pattern from physiological, perceptual, and affective markers. *Prev Med.* 2005 Apr;40(4):407-19.

Lindh-Astrand L, Nedstrand E, Wyon Y, Hammar M. Vasomotor symptoms and quality of life in previously sedentary postmenopausal women randomised to physical activity or estrogen therapy. *Maturitas.* 2004 Jun 15;48(2):97-105.

Lustyk MK, Widman L, Paschane A, Ecker E. Stress, quality of life and physical activity in women with varying degrees of premenstrual symptomatology. *Women Health.* 2004;39(3):35-44.

Lynch AM, McDonald S, Magann EF, Evans SF, Choy PL, Dawson B, et al. Effectiveness and safety of a structured swimming program in previously sedentary women during pregnancy. *J Matern Fetal Neonatal Med* 2003;14:163-9.

Mirzaiinjtabadi K, Anderson D, Barnes M. The relationship between exercise, Body Mass Index and menopausal symptoms in midlife Australian women. *Int J Nurs Pract.* 2006 Feb;12(1):28-34.

Monk C. Stress and mood disorders during pregnancy: implications for child development. *Psychiatr Q.* 2001 Winter;72(4):347-57. Review.

Monk C, Myers MM, Sloan RP, Ellman LM, Fifer WP. Effects of women's stress-elicited physiological activity and chronic anxiety on fetal heart rate. *J Dev Behav Pediatr.* 2003 Feb;24(1):32-8.

Morris SN, Johnson NR. Exercise during pregnancy: a critical appraisal of the literature. *J Reprod Med.* 2005 Mar;50(3):181-8. Review.

Parnpiansil P, Jutapakdeegul N, Chentanez T, Kotchabhakdi N. Exercise during pregnancy increases hippocampal brain-derived neurotrophic factor mRNA expression and spatial learning in neonatal rat pup. *Neurosci Lett*. 2003 Nov 27;352(1):45-8.

Pearlstein T, Steiner M. Non-antidepressant treatment of premenstrual syndrome. *J Clin Psychiatry*. 2000;61 Suppl 12:22-7.

Perrson I, Bergkivist, L, Lindgren, C., Yuen, J. (1997). Hormone replacement therapy and major risk factors for reproductive cancers, osteoporosis, and cardiovascular diseases: evidence of confounding by exposure characteristics. *J. of Clin. Epidemiol.*, 50, 611-618.

Pivarnik JM (Chair) roundtable. Impact of physical activity during pregnancy and postpartum on chronic disease risk. *Med Sci Sports Exerc*. 2006 May;38(5):989-1006.

Polley BA, Wing RR, Sims CJ. Randomized controlled trial to prevent excessive weight gain in pregnant women. *Int J Obes Relat Metab Disord* 2002;26:1494-502.

Poudevigne MS, O'Connor PJ. A review of physical activity patterns in pregnant women and their relationship to psychological health. *Sports Med*. 2006;36(1):19-38. Review.

Poudevigne MS, O'Connor PJ. Physical activity and mood during pregnancy. *Med Sci Sports Exerc*. 2005 Aug;37(8):1374-80.

Poudevigne MS, O'Connor PJ. Physical activity and mood during pregnancy. *Med Sci Sports Exerc*. 2005 Aug;37(8):1374-80.

Poudevigne MS, O'Connor PJ. A review of physical activity patterns in pregnant women and their relationship to psychological health. *Sports Med*. 2006;36(1):19-38. Review.

Redila VA, Olson AK, Swann SE, Mohades G, Webber AJ, Weinberg J, Christie BR. Hippocampal cell proliferation is reduced following prenatal ethanol exposure but can be rescued with voluntary exercise. *Hippocampus*. 2006;16(3):305-11.

Rooks DS, Gautam S, Romeling M, Cross ML, Stratigakis D, Evans B, Goldenberg DL, Iversen MD, Katz JN. Group exercise, education, and combination self-management in women with fibromyalgia: a randomized trial. *Arch Intern Med*. 2007 Nov 12;167(20):2192-20.

Sarsan A, Ardiç F, Ozgen M, Topuz O, Sermez Y. The effects of aerobic and resistance exercises in obese women. *Clin Rehabil*. 2006 Sep;20(9):773-82.

Sheline YI, Sanghavi M, Mintun MA, Gado MH. Depression duration but not age predicts hippocampal volume loss in medically healthy women with recurrent major depression. *J Neurosci*. 1999 Jun 15;19(12):5034-43.

Shangold M, Rebar RW et al (1990) Evaluation and management of menstrual dysfunction in athletes. *Jama* 263(12):1665-1669

Steege JF, Blumenthal JA (1993) The effects of aerobic exercise on premenstrual symptoms in middle-aged women: a preliminary study. *J Psychosom Res* 37(2):127–133

Stoddard JL, Dent CW, Shames L, Bernstein L. Exercise training effects on premenstrual distress and ovarian steroid hormones. *Eur J Appl Physiol*. 2007 Jan;99(1):27-37.

Uzendoski AM, Latin RW, Berg KE, Moshier S. Physiological responses to aerobic exercise during pregnancy and post-partum. *J Sports Med Phys Fitness* 1990;30:77–82.

Watson WJ, Katz VL, Hackney AC, Gall MM, McMurray RG. Fetal responses to maximal swimming and cycling exercise during pregnancy. *Obstet Gynecol* 1991;77:382–6.

Weltman A, Weltman JY, Roy CP, Wideman L, Patrie J, Evans WS, Veldhuis JD. Growth hormone response to graded exercise intensities is attenuated and the gender difference abolished in older adults. *J Appl Physiol*. 2006 May;100(5):1623-9. Epub 2005 Dec 29.

Weuve, J., Kang, J.H., Manson, J.E., Breteler, M.M.B., Ware, J.H. & Grodstein, F. (2004). Physical activity including walking and cognitive function in older women. *Journal of the American Medical Association*, 292, 1454-1461.

Whiting, P., Bagnall, A., Sowden, A., et al (2001) Interventions for the treatment and management of chronic fatigue syndrome: a systematic review. *JAMA*, 286, 1360-1388.

Yaffe, K., Barnes, D., Nevitt, M., Lui, L.Y., Covinsky, K. (2001). A prospective study of physical activity and cognitive decline in elderly women. *Archives of Internal Medicine*, 161, 1703-1708.

Yang J, Li W, Liu X, Li Z, Li H, Yang G, Xu L, Li L. Enriched environment treatment counteracts enhanced addictive and depressive-like behavior induced by prenatal chronic stress. *Brain Res*. 2006 Dec 13;1125(1):132-7.

Yang J, Hou C, Ma N, Liu J, Zhang Y, Zhou J, Xu L, Li L. Enriched environment treatment restores impaired hippocampal synaptic plasticity and cognitive deficits induced by prenatal chronic stress. *Neurobiol Learn Mem*. 2007 Feb;87(2):257-63.

Yang J, Han H, Cao J, Li L, Xu L. Prenatal stress modifies hippocampal synaptic plasticity and spatial learning in young rat offspring. *Hippocampus*. 2006;16(5):431-6.